

Modular Construction of Fixed Point Combinators and Clocked Böhm Trees

Jörg Endrullis Dimitri Hendriks Jan Willem Klop

Vrije Universiteit Amsterdam, Department of Computer Science
De Boelelaan 1081a, 1081 HV Amsterdam, The Netherlands

joerg@few.vu.nl diem@cs.vu.nl jwk@cs.vu.nl

Abstract

Fixed point combinators (and their generalization: looping combinators) are classic notions belonging to the heart of λ -calculus and logic. We start with an exploration of the structure of fixed point combinators (fpc's), vastly generalizing the well-known fact that if Y is an fpc, $Y(SI)$ is again an fpc, generating the Böhm sequence of fpc's. Using the infinitary λ -calculus we devise infinitely many other generation schemes for fpc's. In this way we find schemes and building blocks to construct new fpc's in a modular way.

Having created a plethora of new fixed point combinators, the task is to prove that they are indeed new. That is, we have to prove their β -inconvertibility. Known techniques via Böhm Trees do not apply, because all fpc's have the same Böhm Tree (BT). Therefore, we employ 'clocked BT's', with annotations that convey information of the tempo in which the data in the BT are produced. BT's are thus enriched with an intrinsic clock behaviour, leading to a refined discrimination method for λ -terms. The corresponding equality is strictly intermediate between $=_\beta$ and $=_{BT}$, the equality in the classical models of λ -calculus. An analogous approach pertains to Lévy–Longo and Berarducci trees. Finally, we increase the discrimination power by a precision of the clock notion that we call 'atomic clock'.

The theory of sage birds (technically called *fixed point combinators*) is a fascinating and basic part of combinatory logic; we have only scratched the surface.

R. Smullyan [17].

1 Introduction

Böhm trees constitute a well-known method to discriminate λ -terms M , N : if $BT(M)$ and $BT(N)$ are not identical, then M and N are β -inconvertible, $M \neq_\beta N$. But how do we prove β -inconvertibility of λ -terms with the same BT?

This question was raised in Scott [16] for the interesting equation $BY = BYS$ between terms that as Scott noted are presumably β -inconvertible, yet BT-equal ($=_{BT}$). Scott used his Induction Rule to prove that $BY =_{BT} BYS$; instead we will employ below the infinitary λ -calculus with the same effect, but with more convenience for calculations as a direct generalization of finitary λ -calculus. Often one can solve such a β -discrimination problem by finding a suitable invariant for all the β -reducts of M, N . Below we will do this by way of preparatory example for the fpc's in the Böhm sequence. But a systematic method for this discrimination problem has been lacking, and such a method is one of the two contributions of this paper.




Figure 1: Comparison of (atomic) clock semantics and unlocked semantics. Higher means more identifications.

Actually, the need for such a strategic method was forced upon us, by the other contribution, because Scott's equation $BY = BYS$ turned out to be the key unlocking a plethora of new fpc's. The new generation schemes are of the form: if Y is a fpc, then $YP_1 \dots P_n$ is a fpc, abbreviated as $Y \Rightarrow YP_1 \dots P_n$. So $\square P_1 \dots P_n$ is a 'fpc-generating' vector, and can be considered as a building block to make new fpc's. But are they indeed new? A well-known example of a (singleton)-fpc-generating vector is $\square \delta$, where $\delta = SI$, giving rise when starting from Curry's fpc to the Böhm sequence of fpc's. Here another interesting equation is turning up, namely $Y = Y\delta$, for an arbitrary fpc Y , considered by Statman and Intrigila. In fact, it is implied by Scott's equation:

$$BY = BYS \implies BYI = BYSI \iff Y = Y\delta$$

The first equation $BY = BYS$ will yield many new fpc's, built in a modular way; the last equation $Y = Y\delta$ addresses the question whether they are indeed new. Finding ad hoc invariant proofs for their novelty is too cumbersome. But fortunately, it turns out that although the new fpc's all have the same BT, namely $\lambda f.f^\omega$, they differ in the way this BT is formed, in the 'tempo of formation', where the ticks of the clock are head reduction steps. More generally, we can discern a clock-like behaviour of BT's, that enables us to discriminate the

terms in question. However, this refined discrimination method does not work for all λ -terms; only for a class of ‘simple’ terms, that still is fairly extensive; it includes all fpc’s that are constructed in the modular way that we present, thereby solving our novelty problem. In fact, we gain some more ground: though our discrimination method works best for pairs of simple terms, it can also fruitfully be applied to compare a simple term with a non-simple term, and with some more effort, we can even compare and discriminate two non-simple terms.

Even so, many pairs of fpc’s cannot yet be discriminated, because they not only have the same BT, they also have the same clocked BT. Therefore, in a final grading up of the precision, we introduce ‘atomic clocks’, where the actual position of a head reduction step is administrated. All this pertains not only to the BT-semantics, but also to Lévy–Longo Trees (LLT) (or lazy trees), and Berarducci Trees (BeT) (or syntactic trees). Many problems stay open, in particular problems generalizing the equation of Statman and Intrigila, when arbitrary fpc’s are considered — indeed, we have only scratched the surface.

2 Preliminaries

To make this paper moderately self-contained, and to fix notations, we lay out some ingredients. For λ -calculus we refer to [2] and [5]. For an introduction to Böhm, Berarducci and Lévy–Longo trees, we refer to [2, 1, 6, 3].

Definition 1. λ -terms are defined by the grammar:

$$M ::= x \mid \lambda x.M \mid MM$$

We let $Ter(\lambda)$ denote the set of λ -terms, and use $M, N, \dots$ to range over the elements of $Ter(\lambda)$. The relation $\rightarrow_\beta$ is the compatible closure (i.e., closure under term formation) of the β -rule:

$$(\lambda x.M)N \rightarrow M[N/x] \quad (\beta)$$

where $M[N/x]$ denotes the result of substituting N for all free occurrences of x in M . Furthermore, we use $\twoheadrightarrow_\beta$ to denote the reflexive–transitive closure of $\rightarrow_\beta$. We write $M =_\beta N$ to denote that M is β -convertible with N , i.e., $=_\beta$ is the equivalence closure of $\rightarrow_\beta$. For syntactic equality (modulo renaming of bound variables), we use $\equiv$. We will often omit the subscript β in $\rightarrow_\beta$ and $\twoheadrightarrow_\beta$, but not so for $=_\beta$, in order to reserve $=$ for definitional equality.

A λ -term M are called *normal form* if there exists no N with $M \rightarrow N$. We say that a term M *has a normal form* if it reduces to one. For λ -terms M having a normal form we write $\underline{M}$ for the unique normal form N with $M \rightarrow N$ (note that uniqueness follows from confluence of the λ -calculus).

Some commonly used combinators are:

$$I = \lambda x.x \qquad S = \lambda xyz.xz(yz) \qquad B = \lambda xyz.x(yz)$$

Definition 2. A *position* is a sequence over $\{0, 1, 2\}$. The *subterm* $M|_p$ of M at position p is defined by:

$$\begin{aligned} M|_\epsilon &= M & (MN)|_{1p} &= M|_p \\ (\lambda x.M)|_{0p} &= M|_p & (MN)|_{2p} &= N|_p \end{aligned}$$

$\text{Pos}(M)$ is the set of positions p such that $M|_p$ is defined.

Definition 3.

- (i) A term Y is an *fpc* if $Yx =_\beta x(Yx)$.
- (ii) An fpc Y is *k-reducing* if $Yx \rightarrow^k x(Yx)$.
- (iii) A term Z is a *weak fpc (wfpc)* if $Zx =_\beta x(Z'x)$ where Z' is a wfpc.

A wfpc is alternatively defined as a term having the same Böhm tree as an fpc, namely $\lambda x.x^\omega \equiv \lambda x.x(x(x(\dots)))$.

Weak fpc's are known in foundational studies of type systems as *looping combinators*; see, e.g., [8] and [10].

Example 4. Define by double recursion, Z and Z' such that $Zx = x(Z'x)$ and $Z'x = x(Zx)$. Then Z, Z' are both wfpc's, and $Zx = x(x(Zx))$. So Z delivers its output twice as fast as an ordinary fpc, but the generator flipflops.

As to 'double recursion', [14] collects several proofs of the double fixed point theorem, including some in [2, 17].

Definition 5.

- (i) A *head reduction step* $\rightarrow_h$ is a β -reduction step of the form:

$$\lambda x_1 \dots x_n.(\lambda y.M)NN_1 \dots N_m \rightarrow \lambda x_1 \dots x_n.(M[y/N])N_1 \dots N_m$$

with $n, m \geq 0$.

- (ii) Accordingly, a *head normal form (hnf)* is a λ -term of the form

$$\lambda x_1 \dots \lambda x_n.yN_1 \dots N_m$$

with $n, m \geq 0$.

- (iii) A *weak head normal form (whnf)* is an hnf or an abstraction, that is, a whnf is a term of the form $xM_1 \dots M_m$ or $\lambda x.M$.
- (iv) A term *has a (weak) hnf* if it reduces to one.
- (v) We call a term *root-stable* if it does not reduce to a redex: $(\lambda x.M)N$. A term is called *root-active* if it does not reduce to a root-stable term.

Infinitary λ -calculus $\lambda^\infty\beta$. We will only use the infinitary λ -calculus $\lambda^\infty\beta$ for some simple calculations such as $(\lambda ab.(ab)^\omega)I = \lambda b.(Ib)^\omega = \lambda b.b^\omega$. For a proper setup of $\lambda^\infty\beta$ we refer to [4, 13, 12, 3]. In a nutshell, $\lambda^\infty\beta$ extends finitary λ -calculus by admitting infinite λ -terms, the set of which is called $Ter^\infty(\lambda)$, and infinite reductions (in [19, Ch. 12] and [3] possibly transfinitely long, in [4] of length $\leq \omega$). Limits of infinite reduction sequences are obtained by a strengthening of Cauchy-convergence, stipulating that the depth of contracted redexes must tend to infinity. The $\lambda^\infty\beta$ -calculus is not infinitary confluent (CR^∞), but still has unique infinite normal forms (UN^∞). Böhm Trees (BT's) without $\perp$ are infinite normal forms in $\lambda^\infty\beta$. But beware, the reverse does not hold, e.g. $\lambda x.(\lambda x.(\lambda x.\dots$ is an infinite normal form, but not a BT; it is in fact an LLT (Lévy–Longo Tree, and also a BeT (Berarducci Tree). The notions BT, LLT, BeT are defined e.g. in [3], and in [6]. These notions are also defined in Sections 6 and 8, via their clocked versions.

Definition 6. For terms A, B we define $AB^{\sim n}$ and $A^n B$:

$$\begin{aligned} AB^{\sim 0} &= A & A^0 B &= B \\ AB^{\sim n+1} &= ABB^{\sim n} & A^{n+1} B &= A(A^n B) \end{aligned}$$

A context of the form $\square B^{\sim n}$ is called a *vector*. For the vector notation, it is to be understood that term formation gets highest priority, i.e., $ABC^{\sim n} = (AB)C^{\sim n}$.

3 The Böhm Sequence

There are several ways to make fpc's. For heuristics behind the construction of Curry's fpc $Y_0 = \lambda f.\omega_f\omega_f$, with $\omega_f = \lambda x.f(xx)$, and Turing's fpc $Y_1 = \eta\eta$ with $\eta = \lambda x.f.f(xxf)$, see [14]. The following is an easy exercise.

Proposition 7. *The fpc's Y_0 and Y_1 are not β -convertible.*

It is well-known, as observed by C. Böhm [7] and others, that the class of fpc's coincides exactly with the class of fixed points of the peculiar term $\delta = \lambda ab.b(ab)$, convertible with SI . The notation δ is convenient for calculations and stems from [11].

This term also attracted the attention of R. Smullyan, in his beautiful fable about fpc's figuring as birds in an enchanted forest: “An extremely interesting bird is the owl O defined by the following condition: $Oxy = y(xy)$.” [17]. We will return to the Owl in Remark 10 below.

Thus the term $Y\delta$ is an fpc whenever Y is. It follows that starting with Y_0 , Curry's fpc, we have an infinite sequence of fpc's $Y_0, Y_0\delta, Y_0\delta\delta, \dots, Y_0\delta^{\sim n}, \dots$. We call this sequence the *Böhm sequence*. We will indicate $Y_0\delta^{\sim n}$ by Y_n . Note that indeed Y_1 , the notation that we had given to Turing's fpc, is correct in this naming convention. Now the question is whether all these ‘derived’ fpc's are really new, in other words, whether the sequence is free of duplicates. This is *Exercise 6.8.9 in [2].

Note that we could also have started the sequence from another fpc than Curry's. Now for the sequence starting from an *arbitrary* fpc Y , it is actually an open problem whether that sequence of fpc's $Y, Y\delta, Y\delta\delta, \dots, Y\delta^{\sim n}, \dots$ is free of repetitions. All we know, applying Intrigila's theorem, Theorem 9 below, is that no two consecutive fpc's in this sequence are convertible. But let us first consider the Böhm sequence.

We show that the Böhm sequence contains no duplicates by determining the set of reducts of every Y_n . For $Y_3 \equiv \eta\eta\delta\delta$, the head reduction is displayed in Figure 2, but this is by no means the whole reduction graph. For future




Figure 2: Head reduction of $Y_3 x$.

reference we note that the head reduction diagram suggests a ‘clock behaviour’.

Theorem 8. *The Böhm sequence contains no duplicates.*

Proof. (See also [14].) We define languages $\mathcal{L}_n \subseteq \text{Ter}(\lambda)$ where $\mathcal{L}_n$ is the set of $\rightarrow$ -reducts of Y_n . For $n \geq 1$ we take $Y_n \equiv \eta\eta\delta^{\sim n-1}$, where $\eta \equiv \lambda x f.f(xxf)$.

$$\begin{aligned} \mathcal{L}_0 &::= \lambda f.f^k(\omega_f \omega_f) & (k \geq 0) \\ \mathcal{L}_1 &::= \eta\eta \mid \lambda f.f^k(\mathcal{L}_1 f) & (k > 0) \\ \mathcal{L}_n &::= \mathcal{L}_{n-1}\delta \mid \delta\mathcal{L}_n \mid \lambda b.\delta^k(\mathcal{L}_n b) & (n > 1, k > 0) \end{aligned}$$

Then we show that:

- (i) $\mathcal{L}_n$ is closed under β -reduction; and
- (ii) $\mathcal{L}_n$ and $\mathcal{L}_m$ are disjoint, for $n \neq m$.

This implies that $Y_n \not\equiv_\beta Y_m$ for all $n \neq m$.

For $n \neq m$, $n > 1$, (ii) follows by counting the number of passive δ 's. An occurrence of δ is passive if it occurs as $P\delta$ for some P . To see that $\mathcal{L}_0 \cap \mathcal{L}_1 = \emptyset$, note that if $M \in \mathcal{L}_1$ is an abstraction then $M \equiv \lambda f.f^k(Pf)$ containing a subterm Pf which is never the case in $\mathcal{L}_0$.

We show (i): if $M \in \mathcal{L}_n$ and $M \rightarrow N$, then $N \in \mathcal{L}_n$. Using induction, we do not need to consider cases where the rewrite step is inside a variable of the grammar. We write $\mathcal{L}_n$ in terms as shorthand for a term $M \in \mathcal{L}_n$.

($\mathcal{L}_0$) We have $\lambda f.f^k(\omega_f \omega_f) \rightarrow \lambda f.f^{k+1}(\omega_f \omega_f) \in \mathcal{L}_0$.

($\mathcal{L}_1$) We have $\eta\eta \rightarrow \lambda f.f(\eta\eta f) \in \mathcal{L}_1$,
and $\lambda f.f^k(\lambda f.f^\ell(\mathcal{L}_1 f)f) \rightarrow \lambda f.f^{k+\ell}(\mathcal{L}_1 f) \in \mathcal{L}_1$.

- ($\mathcal{L}_n$) Case 1: $\lambda f.f^k(\mathcal{L}_1 f)\delta \rightarrow \delta^k(\mathcal{L}_1 \delta) \in \mathcal{L}_n$ for $n = 2$, and $(\lambda b.b^k(\mathcal{L}_{n-1} b))\delta \rightarrow \delta^k(\mathcal{L}_{n-1} \delta) \in \mathcal{L}_n$ for $n > 2$.
Case 2: $\delta \mathcal{L}_n \rightarrow \lambda b.b(\mathcal{L}_n b) \in \mathcal{L}_n$.
Case 3: $\lambda b.b^k(\lambda c.c^\ell(\mathcal{L}_n c)b) \rightarrow \lambda b.b^{k+\ell}(\mathcal{L}_n b) \in \mathcal{L}_n$.

□

A very interesting theorem involving δ was proved by B. Intrigila, affirming a conjecture by R. Statman.

Theorem 9 (Intrigila [11]). *There is no ‘double’ fixed point combinator. That is, for no fpc Y we have $Y\delta =_\beta Y$.*

Remark 10 (Smullyan’s Owl $SI =_\beta \delta \equiv \lambda xy.y(xy)$).

We collect some salient facts and questions.

- (i) If Y is an fpc, then $Y\delta$ is an fpc [7].
- (ii) Let Z be a wfpc. Then both δZ and $Z\delta$ are wfpc’s [17].
- (iii) Call an applicative combination of δ ’s a δ -term. In spite of δ ’s simplicity, not all δ -terms are strongly normalizing (SN). An example of a δ -term with infinite reduction is $\delta\delta(\delta\delta)$ (Johannes Waldman, Hans Zantema, personal communication, 2007).
- (iv) Let t be a non-trivial δ -term, i.e., not a single δ . Then t is SN iff t contains exactly one occurrence of $\delta\delta$. Furthermore, if δ -terms t, t' are SN, then they are convertible iff t, t' have the same length [3].
- (v) Convertibility is decidable for δ -terms [18].
- (vi) Call $\Delta = \delta^\omega$, so $\Delta \equiv \delta\Delta$. Then, the infinite λ -term Δ is an fpc: $\Delta x \equiv \delta\Delta x \rightarrow x(\Delta x)$. Δ can be normalized again: $\Delta \rightarrow_\omega \lambda f.f^\omega$. There are many more infinitary fpc’s, e.g. for every n , the infinite term $(SS)^\omega S^{\sim n} I$ is one, as will be clear from the sequel.
- (vii) $\text{BT}(\delta\delta(\delta\delta)) \equiv \perp$, $\delta\delta(\delta\delta)$ has no hnf. Its Berarducci tree is not trivial. Zantema remarked that δ -terms, even infinite ones, such as $\Delta\Delta$, are “top-terminating” (Zantema restricted himself to the applicative rule for δ only — we expect that his observation remains valid for the $\lambda\beta$ -version).
- (viii) Is Intrigila’s theorem also valid for wfpc’s: for no wfpc Z we have $Z\delta =_\beta Z$?

4 The Scott Sequence

In [16] the equation $BY = BYS$ is mentioned as an interesting example of an equation not provable in $\lambda\beta$, while easily provable with Scott's Induction Rule.¹ Scott mentions that he expects that using 'methods of Böhm' the non-convertibility in $\lambda\beta$ can be established, but that he did not attempt a proof. On the other hand, with the induction rule the equality is easily established. We will not consider Scott's Induction Rule, but we will be working in the infinitary lambda calculus, $\lambda^\infty\beta$. It is readily verified that in $\lambda^\infty\beta$ we have:

$$BY = BYS = \lambda ab.(ab)^\omega$$

Proposition 11. $BY_0 \neq_\beta BY_0S$

Proof. Postfixing the combinator I yields BY_0I and BY_0SI . Now $BY_0I =_\beta Y_0$ and $BY_0SI =_\beta Y_0(SI) = Y_1$. Because $Y_0 \neq Y_1$ (Proposition 7), the result follows. $\square$

In the same breath we can strengthen this non-equation to all fpc's Y , by the same calculation followed by an application of Theorem 9 stating that for no fpc Y we have $Y =_\beta Y\delta =_\beta Y(SI)$.

Remark 12.

- (i) The idea of postfixing an I is suggested by the BT $\lambda ab.(ab)^\omega$ of BY and BYS . Namely, in $\lambda^\infty\beta$ we calculate: $(\lambda ab.(ab)^\omega)I = \lambda b.(Ib)^\omega = \lambda b.b^\omega$ which is the BT of any fpc.
- (ii) Interestingly, Scott's equation $BY = BYS$ implies the equation of Statman and Intrigila, $Y = Y\delta$ as one readily verifies, as in the proof of Proposition 11.

Actually, the comparison between the terms BY and BYS has more in store for us than just providing an example that the extension from finitary lambda calculus $\lambda\beta$ to infinitary lambda calculus $\lambda^\infty\beta$ is not conservative. The BT-equality of BY and BYS suggests looking at the whole sequence $BY, BYS, BYSS, BYSSS, \dots, BYS^{\sim n}, \dots$. By the congruence property of BT-equality, they all have the same BT $\lambda ab.(ab)^\omega$; so the terms in this sequence are not fpc's. But they are close to being fpc's, for the first two terms in the sequence we already saw above that postfixing an I turns them into fpc's Y_0, Y_1 . How about postfixing an I to all the terms in the sequence, yielding

$$BYI, BYSI, BYSSI, BYSSSI, \dots, BYS^{\sim n}I, \dots$$

to which we will refer as the *Scott sequence*. Surprisingly, all these terms are fpc's. The Scott sequence concurs with the Böhm sequence of fpc's only for the first two elements, and then splits off with different fpc's. But there is a second surprise. In showing that $BYS^{\sim n}I$ is an fpc, we find as a bonus the fpc-generating vector $\square(SS)S^{\sim n}I$ (which does preserve reducingness). We collect the result.

¹This equation is also discussed in [9].

Theorem 13. *Let Y be a k -reducing fpc. Then:*

- (i) $BYS^{\sim n}I$ is a (non-reducing) fpc, for all $n \geq 0$;
- (ii) $Y(SS)S^{\sim n}I$ is a $(k + 3n + 7)$ -reducing fpc, for all $n \geq 0$.

The proof of Theorem 13 is easy: see the next example.

Example 14. Let Y be a k -reducing fpc. Then:

$$\begin{aligned} BYSSSIx &\rightarrow_h Y(SS)SIx \rightarrow_h^k SS(Y(SS))SIx \\ &\rightarrow_h^3 SS(Y(SS)S)Ix \rightarrow_h^3 SI(Y(SS)SI)x \\ &\rightarrow_h^3 Ix(Y(SS)SIx) \rightarrow_h^1 x(Y(SS)SIx) \end{aligned}$$

This shows that $BYS^{\sim 3}I$ is a non-reducing fpc, and at the same time that $Y(SS)SIx$ is reducing.

Remark 15. Another ‘fpc-generating vector’ is obtained as follows. Start with the equation $Mab = ab(Mab)$; solutions all have the BT seen above, $\lambda ab.(ab)^\omega$. For every M satisfying this equation, we have that MI is an fpc. For: $MIx = Ix(MIx) = x(MIx)$. Now we can solve the equation in different ways.

- (i) $Mab = Y(ab)$, so $M = \lambda ab.Y(ab) = (\lambda yab.y(ab))Y = BY$, as found before.
- (ii) $Mab = ab(Mab) = Sa(Ma)b$, which is obtained by solving $Ma = Sa(Ma)$, leading to $Ma = Y(Sa) = BYSa$, so $M = BYS$. Also this solution was considered before.
- (iii) $M = \lambda ab.ab(Mab) = (\lambda mab.ab(mab))M$, yielding $M = Y\varepsilon$ with $\varepsilon = \lambda abc.bc(abc)$. So if Y is an fpc, then $Y\varepsilon I$ is again an fpc.

5 Generalized Generation Schemes

The schemes mentioned in Theorem 13 and Remark 15(iii) for generating new fixed points from old, are by no means the only ones. There are in fact infinitely many of such schemes. They can be obtained analogously to the ones that we extracted above from the equation $BY = BYS = \lambda ab.(ab)^\omega$, or the equation $Mab = ab(Mab)$. We only treat the case for $n = 3$: consider the equation $Nabc = abc(Nabc)$. Then every solution N is again a ‘pre-fpc’, namely NII is a fpc: $NIIx =_\beta IIX(NIIx) =_\beta x(NIIx)$.

- (i) $Nabc = Y(abc)$, which yields $N = (\lambda yabc.y(abc))Y = (\lambda yabc.BBBYabc)Y$. We obtain $N = BBBY$.
- (ii) $N = Y\xi$ with $\xi = \lambda nabc.abc(nabc)$, yielding the fpc-generating vector $\square\xi II$.

- (iii) $Nabc = abc(Nabc) = S(ab)(Nab)c$. So we take $Nab = S(ab)(Nab)$, which yields $Nab = Y(S(ab)) = BBY(BS)ab$. So $N = BBY(BS)$, and thus we find the equation $BBY = BBY(BS)$, in analogy with the equation $BY =_{\beta} BYS$ above.

Also this equation spawns lots of fpc's as well as fpc-generating vectors. Let's abbreviate BS by A . First one forms the sequence

$$BBY, BBYA, BBYAA, BBYAAA, \dots$$

These terms all have the BT $\lambda abc.ab(ab)^{\omega}$. They are not yet fpc's, but only 'pre-fpc's'. But after postfixing this time $\dots II$ we do again obtain a sequence of fpc's:

$$BBYII, BBY AII, BBY A AII, \dots$$

Again the first two coincide with Y_0, Y_1 , but the series deviates not only from the Böhm sequence but also from the Scott sequence above. As above, the proof that a term in this sequence is indeed a fpc, yields a fpc-generating vector. Thus we find e.g. the following new fpc-generating schemes, which we render in a self-explaining notation:

- (i) $Y \Rightarrow Y(S(AI)I)$
- (ii) $Y \Rightarrow Y(AAA)II$
- (iii) $Y \Rightarrow Y(AII)$
- (iv) $Y \Rightarrow Y(AAI)I$
- (v) $Y \Rightarrow Y(AAA)A^{\sim n}II$

(Note: scheme (iii) came up out of the general search; one may recognize that it is not a new scheme, because the term AII is actually the Owl δ). We can derive many more of these schemes by proceeding with solving the general equation $Na_1a_2\dots a_n = a_1a_2\dots a_n(Na_1a_2\dots a_n)$, bearing in mind the following proposition.

Proposition 16. *If N is a term satisfying*

$$Na_1a_2\dots a_n = a_1a_2\dots a_n(Na_1a_2\dots a_n)$$

then $NI^{\sim(n-1)}$ is an fpc.

We finally mention an fpc-generating scheme with 'dummy parameters'.

- (vi) $Y \Rightarrow YQP_1\dots P_n$ where $P_1, \dots, P_n$ are arbitrary (dummy) terms, and $Q = \lambda y p_1\dots p_n x.x(y p_1\dots p_n x)$.

6 Clock Behaviour of Lambda Terms

As we have seen, there is vast space of fpc's and there are many ways to derive new fpc's. The question is whether all these fpc's are indeed new. So we have to prove that they are not β -convertible.

For the Böhm sequence we did this by an ad hoc argument based on a syntactic invariant; and this method works fine to establish lots of non-equations between the alleged 'new' fpc's that we constructed above. Still, the question remains whether there are not more 'strategic' ways of proving such inequalities.

In this section we propose a more strategic way to discriminate terms with respect to β -conversion. The idea is to extract from a λ -term more than just its BT, but also how the BT was formed; one could say, in what tempo, or in what rhythm. A BT is formed from static pieces of information, but these are rendered in a clock-wise fashion, where the ticks of the internal clock are head reduction steps.

In the sequel we write $[k]t$ for the term t where the root is *annotated with* $k \in \mathbb{N}$. Here, term formation binds stronger than annotation $[k]$. For example $[k]MN$ stands for the term $[k](MN)$ (that is, annotating the (non-displayed) application symbol in-between M and N , in contrast to $([k]M)N$). Moreover, for an annotated term t we use $[t]$ to denote the term obtained from t by dropping all annotations (including annotations of subterms).

Definition 17 (Clocked Böhm trees). Let t be a λ -term. The *clocked Böhm tree* $\text{BT}_{\odot}(t)$ of t is an annotated potentially infinite term defined as follows. If t has no hnf, then define $\text{BT}_{\odot}(t)$ as $\perp$. Otherwise, there is a head reduction $t \rightarrow_h^k \lambda x_1 \dots \lambda x_n. y M_1 \dots M_m$ to hnf. Then we define $\text{BT}_{\odot}(t)$ as the term $[k]\lambda x_1 \dots \lambda x_n. y \text{BT}_{\odot}(M_1) \dots \text{BT}_{\odot}(M_m)$.

The (non-clocked) Böhm tree of a λ -term M can be obtained by dropping the annotations: $\text{BT}(M) = [\text{BT}_{\odot}(M)]$.




Figure 3: Clocked Böhm trees of $Y_0 f$ and $Y_1 f$.

Let us consider the fpc's Y_0 of Curry and Y_1 of Turing. We have $Y_0 \equiv \lambda f. \omega_f \omega_f$ where $\omega_f \equiv \lambda x. f x x$, and

$$\omega_f \omega_f \rightarrow_h^1 f(\omega_f \omega_f)$$

Therefore we obtain

$$\text{BT}_{\odot}(Y_0 f) = [2]f \text{BT}_{\odot}(\omega_f \omega_f) \quad \text{and} \quad \text{BT}_{\odot}(\omega_f \omega_f) = [1]f \text{BT}_{\odot}(\omega_f \omega_f).$$

For $Y_1 \equiv \eta\eta$ where $\eta \equiv \lambda x. \lambda f. f(xxf)$ we get:

$$Y_1 f \equiv \eta\eta f \rightarrow_h^2 f(\eta\eta f)$$

Hence, $\text{BT}_{\odot}(Y_1 f) = [2]f\text{BT}_{\odot}(Y_1 f)$. Figure 3 displays the clocked Böhm trees of $Y_0 f$ (left) and $Y_1 f$ (right).

The following definition captures the well-known Böhm equality of λ -terms.

Definition 18. λ -terms M and N are *BT-equal*, denoted $M =_{\text{BT}} N$, if $\text{BT}(M) \equiv \text{BT}(N)$.

If M and N are not BT-equal then $M \neq_{\beta} N$. More generally, if for some F , $\text{BT}(MF) \neq \text{BT}(NF)$ then $M \neq_{\beta} N$. This method is known as *Böhm-out technique* [2].

Below, we refine this approach by comparing the clocked Böhm trees $\text{BT}_{\odot}(M)$ and $\text{BT}_{\odot}(N)$ instead of the ordinary (non-clocked) Böhm trees. In general, $\text{BT}_{\odot}(M) \neq \text{BT}_{\odot}(N)$ does not always imply that $M \neq_{\beta} N$. Nevertheless, for a large class of λ -terms, called ‘simple’ below, this implication will turn out to be true.

We lift relations over natural numbers to relations over clocked Böhm trees.

Definition 19. Let T_1 and T_2 be clocked Böhm trees with $\text{Pos}(T_1) = \text{Pos}(T_2)$, $R \subseteq \mathbb{N} \times \mathbb{N}$, and $p \in \text{Pos}(T_1)$.

We use $T_1 R_p T_2$ to denote that either both $T_1|_p$ and $T_2|_p$ are not annotated, or both are annotated, and $T_1|_p \equiv [k_1]T'_1$ and $T_2|_p \equiv [k_2]T'_2$ with $k_1 R k_2$. If $T_1 R_p T_2$ for every $p \in \text{Pos}(T_1)$, we write $T_1 R T_2$.

We write $T_1 R_{\exists} T_2$, and say R *holds eventually*, if there exists a depth level $\ell \in \mathbb{N}$ such that $T_1 R_p T_2$ for all positions $p \in \text{Pos}(T_1)$ with $|p| \geq \ell$.

Next, we lift relations over clocked Böhm trees to λ -terms.

Definition 20. Let M, N be λ -terms, and $R \subseteq \mathbb{N} \times \mathbb{N}$.

We write $M R_{\odot} N$ whenever $M =_{\text{BT}} N$, and we have that $\text{BT}_{\odot}(M) R \text{BT}_{\odot}(N)$.

We write $M R_{\odot, \infty} N$ if $M =_{\text{BT}} N$, and for infinitely many $p \in \text{Pos}(\text{BT}_{\odot}(M))$ we have $\text{BT}_{\odot}(M) R_p \text{BT}_{\odot}(N)$.

In case of $M \leq_{\odot} N$ ($M \geq_{\odot} N$) we say that M has a *faster* (*slower*) clock than N .

Proposition 21. *Clocks are accelerated under reduction, that is, $\rightarrow \subseteq \geq_{\odot}$, and slowing down under expansion.*

Proof. We proceed by an elementary diagram construction. Whenever we have co-initial steps $M \rightarrow_h M_1$ and $M \rightarrow_{\odot} M_2$, then by orthogonal projection [19] there exist joining steps $M_1 \rightarrow_{\odot} M'$ and $M_2 \rightarrow_h^{\equiv} M'$. Note that the head step $M \rightarrow_h M_1$ cannot be duplicated, only erased in case of an overlap. This leads to the elementary diagram displayed in Figure 4.

We have $\rightarrow \subseteq \rightarrow_{\odot}^*$. By induction on the length of the rewrite sequence $\rightarrow_{\odot}^*$ it suffices to show that $M \rightarrow_{\odot} N$ implies $M \geq N$. Let $M \rightarrow_{\odot} N$.




Figure 4: Elementary diagram.

If M has no hnf, then the same holds for N , and hence $\text{BT}_{\odot}(M) = \perp = \text{BT}_{\odot}(N)$. Therefore assume that there exists a head rewrite sequence $M \rightarrow_h^k H \equiv \lambda x_1. \dots \lambda x_n. y M_1 \dots M_m$ to hnf. We have

$$\text{BT}_{\odot}(M) \equiv [k] \lambda x_1. \dots \lambda x_n. y \text{BT}_{\odot}(M_1) \dots \text{BT}_{\odot}(M_m)$$

Using the elementary diagram above (k times), we can project $M \rightarrow_{\odot} N$ over $M \rightarrow_h^k H$, and obtain $H \rightarrow_{\odot} H'$, $N \rightarrow_h^{\ell} H' \equiv \lambda x_1. \dots \lambda x_n. y M'_1 \dots M'_m$ with $\ell \leq k$. Then $\text{BT}_{\odot}(N) \equiv [\ell] \lambda x_1. \dots \lambda x_n. y \text{BT}_{\odot}(M'_1) \dots \text{BT}_{\odot}(M'_m)$ and $\ell \leq k$. Since $H \rightarrow_{\odot} H'$ and H is in hnf, we get $M_i \rightarrow_{\odot} M'_i$ for every $i = 1, \dots, m$. Co-recursively applying the same argument to $M_i \rightarrow_{\odot} M'_i$ yields $\text{BT}_{\odot}(M) \geq \text{BT}_{\odot}(N)$. $\square$

While $\text{BT}_{\odot}(M) \neq \text{BT}_{\odot}(N)$ does not imply $M \neq_{\beta} N$, the following theorem allows us to use clocked Böhm trees for discriminating λ -terms:

Theorem 22. *Let M and N be λ -terms. Assume there exists a reduct N' of N such that for no reduct M' of M we have $M' \leq_{\odot} N'$. Then $M \neq_{\beta} N$.*

Proof. If $M =_{\beta} N$ then $M =_{\beta} N'$ and $M \twoheadrightarrow M' \leftarrow N'$ for some M' . Hence $M' \leq_{\odot} N'$ by Proposition 21. $\square$

The theorem allows us to pick N' while having to show that $M' \not\leq_{\odot} N'$ for all reducts M' of M . The latter condition is in general difficult to prove. However, the theorem is of use if one of the terms has a manageable set of reducts, and this term happens to have slower clocks.

For a large class of λ -terms it turns out that clocks are invariant under reduction. We call these terms ‘simple’.

Definition 23. A redex $(\lambda x. A)B$ is called:

- (i) *linear* if x has at most one occurrence in A ;
- (ii) *call-by-value* if B is a normal form; and
- (iii) *simple* if it is linear or call-by-value.

The definition of simple redexes generalizes the well-known notions of call-by-value and linear redexes. Next, we define simple *terms*. Intuitively, we call a term t ‘simple’ if every reduction admitted by t only contracts simple redexes. The following definition further generalises this intuition by considering only standard reductions (to normal form):

Definition 24 (Simple terms). A λ -term t is *simple* if either t has no hnf, or the head reduction to hnf $t \rightarrow_h^k \lambda x_1 \dots \lambda x_n. y M_1 \dots M_m$ contracts only simple redexes, and $M_1, \dots, M_m$ are simple terms.

All the fpc's in this paper are either simple or have simple reducts. The clock of simple λ -terms is invariant under reduction, that is, when ignoring finite prefixes of the clocked Böhm trees (by reducing a term we can always make the clock values in a finite prefix equal to 0).

Proposition 25. *Let M, N be λ -terms such that M is a simple term and $M \rightarrow N$. Then $M =_{\rightarrow, \exists}^{\circ} N$, that is, the clocks of M and N are eventually equal.*

Proof. The proof is a straightforward extension of the proof of Proposition 21 with the observation that for simple terms M , rewriting M to hnf:

$$M \rightarrow_h^k H \equiv \lambda x_1 \dots \lambda x_n. y M_1 \dots M_m$$

does not duplicate redexes. Hence, the elementary diagrams are now of the form displayed in Figure 5.




Figure 5: Elementary diagrams for simple M .

That is, whenever we have co-initial steps $M \rightarrow_h M_1$ and $M \rightarrow M_2$ and M is a simple term, then either the steps cancel each other out $M_1 \equiv M_2$ (if both are the same step), or they can be joined by single steps $M_1 \rightarrow_h M' \leftarrow M_2$.

As a consequence, when projecting $M \rightarrow_h M'$ over a rewrite sequence $M \rightarrow^n N$ then either $N \rightarrow^n M'' \leftarrow_h M'$ or there has been cancellation and $N \rightarrow^{n-1} M'' \equiv M'$. Every cancellation decreases the number of steps $N \rightarrow^{n-1} M''$, and hence there can only finitely many cancellations. This implies the claim that $\text{BT}_{\rightarrow, \exists}^{\circ}(M)$ is equal to $\text{BT}_{\rightarrow, \exists}^{\circ}(N)$ modulo a finite prefix, that is, $M =_{\rightarrow, \exists}^{\circ} N$. $\square$

Reduction accelerates clocks, i.e., $\rightarrow \subseteq \geq_{\rightarrow}^{\circ}$. Moreover, for simple terms the clock is invariant under reduction, see Proposition 25. Hence if a term M has a simple reduct N , then N has the fastest clock reachable from M modulo a finite prefix. This justifies the following convention.

Convention 26. The (*minimal*) *clock* of a λ -term M with a simple reduct N is $\text{BT}_{\rightarrow}^{\circ}(N)$, the clocked BT of N .

For simple terms we obtain the following theorem:

Theorem 27. *Let M and N be λ -terms. If there exists a reduct N' of N and a simple reduct M' of M such that $M' >_{\rightarrow, \infty}^{\circ} N'$, then $M \neq_{\beta} N$.*

Proof. Assume $M =_\beta N$. Then $M' = N'$ and by confluence they have a common reduct $M' \rightarrow M'' \leftarrow N'$. Since M' is simple, we have $M' =_{\circlearrowleft, \exists} M''$ by Proposition 25. Hence, $M' >_{\circlearrowleft, \infty} N'$ implies $M'' >_{\circlearrowleft, \infty} N'$, which would contradict $M'' \leq_{\circlearrowleft} N'$ obtained by Proposition 21. $\square$

Theorem 27 significantly reduces the proof obligation in comparison to Theorem 22. We can pick any simple reduct M' of M , instead of having to reason about all reducts M' of M . For the case that both M and N are simple, there is no need to look for reducts:

Proposition 28. *For simple terms M and N , $M \neq_{\circlearrowleft, \infty} N$ implies $M \neq_\beta N$.*

Proof. Assume $M = N$ then $M \rightarrow O \leftarrow N$ for a common reduct O . Then $M =_{\circlearrowleft, \exists} O =_{\circlearrowleft, \exists} N$ by Proposition 25. Hence $M =_{\circlearrowleft, \exists} N$ which contradicts $M \neq_{\circlearrowleft, \infty} N$. $\square$

Example 29. Let $n \geq 2$. We compute the clocks of the fpc's Y_n of the Böhm sequence. We first reduce $Y_n = Y_0 \delta^{\sim n}$ with $Y_0 = \lambda f. \omega_f \omega_f$ and $\omega_f = \lambda x. f(xx)$ to a simple term:

$$Y_n x \rightarrow \omega_\delta \omega_\delta \delta^{\sim n-1} x \rightarrow \underline{\omega_\delta \omega_\delta} \delta^{\sim n-1} x$$

where $\underline{\omega_\delta} = \lambda ab. b(aab)$. We compute the clock:

$$\begin{aligned} \underline{\omega_\delta \omega_\delta} \delta^{\sim n-1} x &\rightarrow_h^2 \delta(\underline{\omega_\delta \omega_\delta} \delta) \delta^{\sim n-2} x \\ &\rightarrow_h^{2(n-2)} \delta(\underline{\omega_\delta \omega_\delta} \delta^{\sim n-1}) x \\ &\rightarrow_h^2 x(\underline{\omega_\delta \omega_\delta} \delta^{\sim n-1} x) \end{aligned}$$

We find $\text{BT}_{\circlearrowleft}(\underline{\omega_\delta \omega_\delta} \delta^{\sim n-1} x) = [2n](x \text{BT}_{\circlearrowleft}(\underline{\omega_\delta \omega_\delta} \delta^{\sim n-1} x))$. Hence, for $n \geq 2$ the clock of Y_n is $2n$.

By Theorem 27, Example 29 and Figure 3 we obtain an alternative proof for Theorem 8: the Böhm sequence contains no duplicates.

Example 30. Let $n \geq 2$. We compute the clocks of the fpc's $U_n = BYS^{\sim n} I$ of the Scott sequence; so where $Y = Y_0$. We first reduce U_n to a simple term:

$$\begin{aligned} U_n x &\rightarrow Y(SS)S^{\sim(n-2)} Ix \\ &\rightarrow \omega_{SS} \omega_{SS} S^{\sim(n-2)} Ix \\ &\rightarrow \underline{\omega_{SS} \omega_{SS}} S^{\sim(n-2)} Ix \end{aligned}$$

where $\underline{\omega_{SS}} = \lambda abc. bc(aabc)$. We abbreviate $\theta \equiv \underline{\omega_{SS}}$. Then we compute the

clocks for $n = 2$, $n = 3$, and $n > 3$:

$$\begin{aligned}
\theta\theta Ix &\rightarrow_h^3 Ix(\theta\theta Ix) \rightarrow_h^1 x(\theta\theta Ix) \\
\theta\theta SIx &\rightarrow_h^3 SI(\theta\theta SI)x \rightarrow_h^4 x(\theta\theta SIx) \\
\theta\theta S^{\sim(n-2)}Ix &\rightarrow_h^3 SS(\theta\theta SS)S^{\sim(n-4)}Ix \\
&\rightarrow_h^{3(n-4)} SS(\theta\theta SSS^{\sim(n-4)})Ix \\
&\rightarrow_h^3 SI(\theta\theta S^{\sim(n-2)}I)x \\
&\rightarrow_h^4 x(\theta\theta S^{\sim(n-2)}Ix)
\end{aligned}$$

respectively. For all three cases, we find:

$$\text{BT}_{\text{c}}(\theta\theta S^{\sim(n-2)}Ix) = [3(n-2)](x\text{BT}_{\text{c}}(\theta\theta S^{\sim(n-2)}Ix))$$

Using Theorem 27 we infer from Example 30 and Figure 3 (recall that $U_0 =_\beta Y_0$ and $U_1 =_\beta Y_1$):

Corollary 31. *The Scott sequence contains no duplicates.*

Plotkin [15] asked: Is there an fpc Y such that

$$A_Y \equiv Y(\lambda z.fzz) =_\beta Y(\lambda x.Y(\lambda y.fxy)) \equiv B_Y$$

or in other notation: $\mu z.fzz =_\beta \mu z.fzz$, where $\mu x.M(x) = Y(\lambda x.M(x))$. The terms A_Y and B_Y have the same Böhm tree, which is the solution of $T = fTT$.

The terms A_Y and B_Y are not simple. An extension of our clock method can be given which restricts the clock comparison to single paths in the clocked Böhm tree along which there is no duplication of redexes. We leave this extension to future work. Using this extension would allow us to settle the question in the negative for all simple fpc's. For Turing's fpc Y_1 this is seen by computing the clocked BT's of A_{Y_1} and B_{Y_1} . Recall $Y_1 \equiv \eta\eta$ with $\eta \equiv \lambda x.f(xxf)$.

$$\begin{aligned}
A_{Y_1} &\equiv \eta\eta(\lambda z.fzz) \rightarrow_h^2 (\lambda z.fzz)A_{Y_1} \\
&\rightarrow_h^1 fA_{Y_1}A_{Y_1} \\
B_{Y_1} &\equiv \eta\eta(\lambda x.\eta\eta(\lambda y.fxy)) \\
&\rightarrow_h^2 (\lambda x.\eta\eta(\lambda y.fxy))B_{Y_1} \\
&\rightarrow_h^1 \eta\eta(\lambda y.fB_{Y_1}y) \\
&\rightarrow_h^2 (\lambda y.fB_{Y_1}y)(\eta\eta(\lambda y.fB_{Y_1}y)) \\
&\rightarrow_h^1 fB_{Y_1}(\eta\eta(\lambda y.fB_{Y_1}y))
\end{aligned}$$

Note that for B_{Y_1} developing the left branch takes six steps, whereas the right only needs three. The clocked BT's for A_{Y_1} and B_{Y_1} are depicted in Figure 6 using hnf-notation (see [2] or [3]).

We conjecture that for no fpc Y , $A_Y =_\beta B_Y$; maybe this requires an extension of the proof in [11].




Figure 6: Clocked BT's for A_{Y_1} and B_{Y_1} .

7 Atomic Clocks

We have introduced clocked Böhm trees for discriminating λ -terms. In this section, we refine the clocks to measure not only the *number* of head steps, but, in addition, the *position* of each of these steps. We call these clocks ‘atomic’.

Let us consider a motivating example. We discriminate Y_2 and U_2 . First, we reduce both terms to simple reducts:

$$\begin{aligned} Y_2 x &\equiv Y_0 \delta \delta x \rightarrow \xi \xi \delta x & \text{where } \xi &= \lambda ab.b(aab) \\ U_2 x &\equiv Y_0 (SS)Ix \rightarrow \theta \theta Ix & \text{where } \theta &= \lambda abc.bc(aabc) \end{aligned}$$

We compute the atomic clocks of these simple reducts:

$$\begin{aligned} \xi \xi \delta x &\rightarrow_{h,11} (\lambda b.b(\xi \xi b))\delta x \rightarrow_{h,1} \delta(\xi \xi \delta)x \\ &\rightarrow_{h,1} (\lambda b.b(\xi \xi \delta b))x \rightarrow_{h,\epsilon} x(\xi \xi \delta x) \\ \theta \theta Ix &\rightarrow_{h,11} (\lambda bc.bc(\theta \theta bc))Ix \rightarrow_{h,1} (\lambda c.Ic(\theta \theta Ic))x \\ &\rightarrow_{h,\epsilon} Ix(\theta \theta Ix) \rightarrow_{h,1} x(\theta \theta Ix) \end{aligned}$$

Both terms have the clocked Böhm tree $T \equiv [4](xT)$. Thus the method from the previous section is not applicable.

However, with atomic clocks we have:

$$\begin{aligned} \text{BT}_{\blacktriangleleft\blacktriangleright}(\xi \xi \delta x) &= [11, 1, 1, \epsilon](x\text{BT}_{\blacktriangleleft\blacktriangleright}(\xi \xi \delta x)) \\ \text{BT}_{\blacktriangleleft\blacktriangleright}(\theta \theta Ix) &= [11, 1, \epsilon, 1](x\text{BT}_{\blacktriangleleft\blacktriangleright}(\theta \theta Ix)) \end{aligned}$$

which allows us to discriminate the terms. Hence $Y_2 \neq U_2$ (by Corollary 28 which generalises to the setting of atomic BT's). Note that the (non-atomic) clocked BT's can be obtained by taking the length of the lists of positions.

For lists $\vec{p}, \vec{q}$ of positions, we write $\vec{p} \cdot \vec{q}$ for concatenating $\vec{p}$ to $\vec{q}$. We write $\rightarrow_{h, \langle p_1, \dots, p_n \rangle}$ for the rewrite sequence $\rightarrow_{h, p_1} \cdots \rightarrow_{h, p_n}$ consisting of steps at position $p_1, \dots, p_n$.

Definition 32 (Atomic clock Böhm trees). Let $t \in \text{Ter}(\lambda)$. The *atomic clock Böhm tree* $\text{BT}_{\blacktriangleleft\blacktriangleright}(t)$ of t is an annotated infinite term defined as follows. If t has no hnf, then define $\text{BT}_{\blacktriangleleft\blacktriangleright}(t)$ as $\perp$. Otherwise, there is a head reduction

$$t \rightarrow_{h, p_1} \cdots \rightarrow_{h, p_k} \lambda x_1 \dots \lambda x_n. y M_1 \dots M_m$$

of length k to hnf. Then we define

$$\text{BT}_{\bullet, \bullet}(t) = [\langle p_1, \dots, p_k \rangle] \lambda x_1. \dots \lambda x_n. y \text{BT}_{\bullet, \bullet}(M_1) \dots \text{BT}_{\bullet, \bullet}(M_m)$$

The theory developed for (non-atomic) BT's generalises as follows to the atomic trees. For lists of positions $\vec{p}, \vec{q}$ we define $\vec{p} \geq \vec{q}$ whenever $\vec{q}$ is a subsequence of $\vec{p}$, and $\vec{p} > \vec{q}$ if additionally $\vec{p} \neq \vec{q}$. Here $\langle a_1, \dots, a_n \rangle$ is a subsequence of $\langle b_1, \dots, b_m \rangle$ if there exist indexes $i_1 < i_2 < \dots < i_n$ such that $\langle a_1, \dots, a_n \rangle = \langle b_{i_1}, \dots, b_{i_n} \rangle$.

Using this notation for comparing the atomic annotations (lists of positions), Proposition 21, Theorem 22, Proposition 25, Theorem 27, and Corollary 28 remain valid (using basically the same proofs).

Proposition 33. *Let $B_n = \square(SS)S^{\sim n}I$ the fpc-generating vectors from Theorem 13. For $n_1, \dots, n_k \in \mathbb{N}$ we define $Y^{\langle n_1, \dots, n_k \rangle} = Y_0 B_{n_1} \dots B_{n_k}$. We prove that all these fpc's are inconvertible, that is, $\vec{n} \neq \vec{m}$ implies $Y^{\vec{n}} \neq_\beta Y^{\vec{m}}$.*

This proposition cannot be proved using (non-atomic) clocks, as for example: $\text{BT}_{\bullet, \bullet}(Y^{\langle n, m \rangle}) = \text{BT}_{\bullet, \bullet}(Y^{\langle m, n \rangle})$. We introduce some auxiliary notations. Let $B'_n = \square S^{\sim n}I$, and define $\underline{B}_n = \square(\underline{SS})S^{\sim n}I$ where $\underline{SS} = \lambda abc. bc(abc)$. For $\vec{p} = \langle 1^{m_1}, \dots, 1^{m_k} \rangle$ a list of positions define $\vec{p} \times n = \vec{p} \cdot (\langle 1^{m_1-1}, \dots, 1^{m_k-1} \rangle \times (n-1))$ for $n > 1$ and $\vec{p} \times 1 = \vec{p}$.

Proof. Let $n_1, \dots, n_k \in \mathbb{N}$ and $Y \equiv \theta \theta B'_{n_1} \underline{B}_{n_2} \dots \underline{B}_{n_k}$ where $\theta = \lambda abc. bc(aabc)$, then $Y^{\langle n_1, \dots, n_k \rangle} x \twoheadrightarrow Yx$ (for $k \geq 1$) where Yx is a simple term. Apart from the initial and final steps, the rewrite sequence $Yx \twoheadrightarrow_h x(Yx)$ is composed of k subsequences of the form:

$$\begin{aligned} SI(\theta \theta \dots) \underline{B}_n V_m &\equiv SI(\theta \theta \dots) (\underline{SS}) S^{\sim n} I V_m \\ &\rightarrow_{h, 1^{n+m+3} \times 3} I(\underline{SS})(\theta \theta \dots (\underline{SS})) S^{\sim n} I V_m \\ &\rightarrow_{h, 1^{n+m+2}} \underline{SS}(\theta \theta \dots (\underline{SS})) S^{\sim n} I V_m \\ &\rightarrow_{h, 1^{n+m+1} \times 3} \underline{SS}(\theta \theta \dots (\underline{SS})) S^{\sim n-2} I V_m \\ &\rightarrow_{h, (1^{n+m} \times 3) \times (n-1)} SI(\theta \theta \dots \underline{B}_n) V_m \end{aligned}$$

for every B_n with $n \geq 2$, and vector V_m of length m .

For every B_n there is exactly one occurrence of four consecutive steps at 'decreasing' positions $1^{n+m+2}, 1^{n+m+1}, 1^{n+m}, 1^{n+m-1}$ (btw, this also holds for $n < 2$). Hence, from the distance between these occurrences we can reconstruct the vector $\langle n_1, \dots, n_k \rangle$. This shows that $\vec{n} \neq \vec{m}$ implies that $\text{BT}_{\bullet, \bullet}(Y^{\vec{n}}) =_\exists \text{BT}_{\bullet, \bullet}(Y^{\vec{m}})$ is false, and hence we conclude $Y^{\vec{n}} \neq_\beta Y^{\vec{m}}$ by Corollary 28. $\square$

8 Clocked Lévy–Longo and Berarducci Trees

In fact, there are three main semantics for the λ -calculus: BT, LLT, and BeT (see [1, 4, 6, 12, 3]). The notions from the previous section generalize directly to LLT and BeT semantics.

Definition 34 (Clocked Lévy–Longo trees). Let t be a λ -term. The *clocked Lévy–Longo tree* $\text{LLT}_{\odot}(t)$ of t is an annotated potentially infinite term defined as follows. If t has no whnf, then define $\text{LLT}_{\odot}(t)$ as $\perp$. Otherwise, there exists a head rewrite sequence $t \rightarrow_h^k \lambda x.M$ or $t \rightarrow_h^k xM_1 \dots M_m$ to whnf. In this case, we define $\text{LLT}_{\odot}(t)$ as $[k]\lambda x.\text{LLT}_{\odot}(M)$ or $[k]x\text{LLT}_{\odot}(M_1) \dots \text{LLT}_{\odot}(M_m)$, respectively.

Definition 35 (Clocked Berarducci trees). Let t be a λ -term. The *clocked Berarducci tree* $\text{BeT}_{\odot}(t)$ of t is an annotated potentially infinite term defined as follows. If t is root-active, let $\text{BeT}_{\odot}(t) \equiv \perp$. If $t \rightarrow_h^k s$ rewrites to a root-stable term $s \equiv x$, $s \equiv \lambda x.M$ or $s \equiv MN$, then define $\text{BeT}_{\odot}(t)$ as $[k]x$, $[k]\lambda x.\text{BeT}_{\odot}(M)$ or $[k]\text{BeT}_{\odot}(M)\text{BeT}_{\odot}(N)$, respectively.

Example 36. We consider the terms $A \equiv aa$ with $a \equiv \lambda x.\lambda y.xx$ and $B \equiv bb$ with $b \equiv \lambda x.\lambda y.\lambda z.xx$. We have:

$$\begin{aligned}\text{LLT}_{\odot}(A) &\equiv [1]\lambda y.\text{LLT}_{\odot}(A) \\ \text{LLT}_{\odot}(B) &\equiv [1]\lambda y.\lambda z.\text{LLT}_{\odot}(B)\end{aligned}$$

Thus, in $\text{LLT}_{\odot}(A)$ every λ requires one head reduction step whereas in $\text{LLT}_{\odot}(B)$ every second λ is obtained for ‘free’ (that is, in 0 steps).

We remark that A and B cannot be distinguished in the Böhm tree semantics since $\text{BT}_{\odot}(A) \equiv \text{BT}_{\odot}(B) \equiv \perp$. In the Böhm tree semantics, a term is meaningful only if it has a hnf. The Lévy–Longo semantics weakens this condition to whnf’s, and thereby allows more terms to be distinguished. The Berarducci tree semantics is a further weakening where only root-active terms are discarded as meaningless.

9 Concluding remarks

We conclude with an encompassing conjecture.

Conjecture 37. *Building fpc’s with fcp-generating vectors is a free construction, that is, there are no non-trivial identifications.*

A first step is found in Intrigila’s theorem $Y\delta \neq_{\beta} Y$, for any fpc Y . A second step is that the Böhm sequence is duplicate-free. A third step is found in our proof that the Scott sequence is duplicate-free, and Proposition 33, which states that there are no identifications when starting the construction with Y_0 .

Other parts of the conjecture are as follows. Let Y, Y' fpc’s and $B_1 \dots B_n, C_1 \dots C_k$ be fcp-generating vectors.

- (i) $Y\delta =_{\beta} Y'\delta$ iff $Y =_{\beta} Y'$.
- (ii) $YB_1 \dots B_n =_{\beta} Y'B_1 \dots B_n$ iff $Y = Y'$.
- (iii) $YB_1 \dots B_n \neq_{\beta} YC_1 \dots C_k$ if $B_1 \dots B_n \neq C_1 \dots C_k$.

For general fpc's Y , Y' these conjectures may be beyond current techniques, but for the well-known fpc's of Curry and Turing, and the fpc-generating vectors introduced here, including their versions for $n > 3$, these problems are tractable.

References

- [1] Samson Abramsky and C.-H. Luke Ong. Full Abstraction in the Lazy Lambda Calculus. *Inf. Comput.*, 105(2):159–267, 1993.
- [2] H. P. Barendregt. *The Lambda Calculus. Its Syntax and Semantics*, volume 103 of *Studies in Logic and The Foundations of Mathematics*. North-Holland, revised edition, 1984.
- [3] H. P. Barendregt and J. W. Klop. Applications of Infinitary Lambda Calculus. *Inf. Comput.*, 207(5):559–582, 2009.
- [4] A. Berarducci and B. Intrigila. Church–Rosser λ -theories, Infinite λ -calculus and Consistency Problems. *Logic: From Foundations to Applications*, pages 33–58, 1996.
- [5] I. Bethke. *Lambda Calculus*. Chapter 10 in [19].
- [6] Inge Bethke, Jan Willem Klop, and Roel C. de Vrijer. Descendants and Origins in Term Rewriting. *Inf. Comput.*, 159(1–2):59–124, 2000.
- [7] C. Böhm. The CUCH as a Formal and Description Language. *Annual Review in Automatic Programming*, 3:179–197, 1963.
- [8] Th. Coquand and H. Herbelin. A -Translation and Looping Combinators in Pure Type Systems. *J. Funct. Program.*, 4(1):77–88, 1994.
- [9] M. Dezani-Ciancaglini, P. Severi, and F. J. de Vries. Infinitary Lambda Calculus and Discrimination of Berarducci Trees. *Theor. Comput. Sci.*, 2(298):275–302, 2003.
- [10] H. Geuvers and B. Werner. On the Church–Rosser Property for Expressive Type Systems and its Consequences for their Metatheoretic Study. In *LICS*, pages 320–329, 1994.
- [11] B. Intrigila. Non-Existent Statman’s Double Fixed Point Combinator Does Not Exist, Indeed. *Inf. Comput.*, 137(1):35–40, 1997.
- [12] R. Kennaway and F.-J. de Vries. *Infinitary Rewriting*. Chapter 12 in [19].
- [13] Richard Kennaway, Jan Willem Klop, M. Ronan Sleep, and Fer-Jan de Vries. Infinitary Lambda Calculus. *Theor. Comput. Sci.*, 175(1):93–125, 1997.

- [14] J. W. Klop. New Fixed Point Combinators from Old. In E Barendsen, V. Capretta, H. Geuvers, and M. Niqui, editors, *Reflections on Type Theory, λ -Calculus, and the Mind. Essays Dedicated to Henk Barendregt on the Occasion of his 60th Birthday*, pages 197–210. 2007. Online version: <http://www.cs.ru.nl/barendregt60>.
- [15] G. Plotkin, 2007. Personal communication at the symposium for H. Barendregt’s 60th birthday.
- [16] D. S. Scott. Some Philosophical Issues Concerning Theories of Combinators. In C. Böhm, editor, *Lambda Calculus and Computer Science Theory*, volume 37 of *LNCS*, pages 346–366, 1975.
- [17] R. Smullyan. *To Mock a Mockingbird, and Other Logic Puzzles*. Alfred A. Knopf, New York, 1985.
- [18] R. Statman. The Word problem for Smullyan’s Lark Combinator is Decidable. *Journal of Symbolic Computation*, 7:103–112, 1989.
- [19] Terese. *Term Rewriting Systems*, volume 55 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2003.